

## **Product datasheet for TP700272**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD1C (NM\_001765) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of human CD1C molecule (CD1C), with C-terminal Fc tag,

expressed in human cells, 20 µg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** 

or AA Sequence:

A DNA sequence from TrueORF clone, RC218490, encoding the region (Asn18 - Met302) of

human CD1C

Tag: C-Fc

**Predicted MW:** 59 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** PBS, pH 7.4, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001756

Locus ID: 911

 UniProt ID:
 P29017

RefSeq Size: 2592

**Cytogenetics:** 1q23.1

RefSeq ORF: 999

Synonyms: BDCA1; CD1; CD1A; R7





**Summary:** 

This gene encodes a member of the CD1 family of transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene is broadly distributed throughout the endocytic system via a tyrosine-based motif in the cytoplasmic tail. Alternatively spliced transcript variants of this gene have been observed, but their full-length nature is not known. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome, Transmembrane

Protein Pathways: Hematopoietic cell lineage

## **Product images:**

