

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP701224-B

Thyroglobulin (TG) (NM_003235) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human thyroglobulin (TG), Asn20-End, with C-terminal His and site-specific Biotinylation on the C-terminal AVI-plus tag, expressed in HEK293 cells, 50ug
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	A DNA sequence from TrueORF clone, RC216216, encoding the region(Asn20-End) of TG
Tag:	C-AVI-PLUS
Predicted MW:	306.0 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	PBS, pH 7.4, 10% glycerol
Bioactivity:	The biotin to protein ratio is 1 as determined by Streptavidin Pull-Down Assay.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP 003226</u>
Locus ID:	7038
UniProt ID:	<u>P01266</u>
RefSeq Size:	8453
Cytogenetics:	8q24.22
RefSeq ORF:	8304
Synonyms:	AITD3; TGN




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Thyroglobulin (TG) (NM_003235) Human Recombinant Protein – TP701224-B
Summary:	Thyroglobulin (Tg) is a glycoprotein homodimer produced predominantly by the thryroid gland. It acts as a substrate for the synthesis of thyroxine and triiodothyronine as well as the storage of the inactive forms of thyroid hormone and iodine. Thyroglobulin is secreted from the endoplasmic reticulum to its site of iodination, and subsequent thyroxine biosynthesis, in the follicular lumen. Mutations in this gene cause thyroid dyshormonogenesis, manifested as goiter, and are associated with moderate to severe congenital hypothyroidism. Polymorphisms in this gene are associated with susceptibility to autoimmune thyroid diseases (AITD) such as Graves disease and Hashimoto thryoiditis. [provided by RefSeq, Nov 2009]
Protein Familie	s: Druggable Genome

Protein Pathways:

Product images:



Autoimmune thyroid disease

Purified recombinant protein TG was analyzed by SDS-PAGE gel and Coomossie Blue Staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US