

Product datasheet for **TP710046**

Isocitrate dehydrogenase (IDH1) (NM_005896) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of mutant(R132S) of human isocitrate dehydrogenase 1 (NADP+), soluble (IDH1),with C-terminal DDK tag,expressed in sf9 cells.
Species:	Human
Expression Host:	Sf9
Expression cDNA Clone or AA Sequence:	A DNA sequence from TrueORF clone, RC400099, encoding human full-length Mutant IDH1(R132S)
Tag:	C-DDK
Predicted MW:	47 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_005887
Locus ID:	3417
UniProt ID:	O75874
RefSeq Size:	2339
Cytogenetics:	2q34
RefSeq ORF:	1242
Synonyms:	HEL-216; HEL-S-26; IDCD; IDH; IDP; IDPC; PICD



[View online »](#)

Summary:

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Sep 2013]

Protein Pathways:

Citrate cycle (TCA cycle), Glutathione metabolism, Metabolic pathways

Product images: