

## **Product datasheet for TP710162**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Eph receptor B4 (EPHB4) (NM\_004444) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human EPH receptor B4 (EPHB4), residues 16-539aa, with C-

terminal DDK tag, expressed in sf9, 20ug

Species: Human

**Expression Host:** Sf9

**Expression cDNA Clone** 

or AA Sequence:

A DNA sequence from TrueORF clone, RC208559, encoding the region(Met-Leu16-Ala539) of

Homo sapiens EPHB4

Tag: C-DDK

**Predicted MW:** 57 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 50 mM Tris-HCl, 100 mM glycine, pH 8.0, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 004435

**Locus ID:** 2050

UniProt ID: <u>P54760</u>, <u>Q96L35</u>, <u>Q541P7</u>

RefSeq Size: 4369 Cytogenetics: 7q22.1 RefSeq ORF: 2961

Synonyms: CMAVM2; HFASD; HTK; LMPHM7; MYK1; TYRO11





**Summary:** 

Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The protein encoded by this gene binds to ephrin-B2 and plays an essential role in vascular development. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome, Protein Kinase, Transmembrane

**Protein Pathways:** Axon guidance

## **Product images:**

