

Product datasheet for TP710202

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PTP1B (PTPN1) (NM_002827) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human protein tyrosine phosphatase, non-receptor type 1

(PTPN1), full length, with C-terminal DDK tag, expressed in sf9, 20ug

Species: Human

Expression Host: Sf9

Expression cDNA Clone

or AA Sequence:

A DNA sequence from TrueORF clone, RC204902, encoding human full-length PTP1B

Tag: C-DDK

Predicted MW: 49.8 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 50 mM Tris-HCl, 100 mM glycine, pH 8.0, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002818

Locus ID: 5770

UniProt ID: <u>P18031</u>, <u>A8K3M3</u>

RefSeq Size: 3573

Cytogenetics: 20q13.13

RefSeq ORF: 1305

Synonyms: PTP1B





Summary:

The protein encoded by this gene is the founding member of the protein tyrosine phosphatase (PTP) family, which was isolated and identified based on its enzymatic activity and amino acid sequence. PTPs catalyze the hydrolysis of the phosphate monoesters specifically on tyrosine residues. Members of the PTP family share a highly conserved catalytic motif, which is essential for the catalytic activity. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP has been shown to act as a negative regulator of insulin signaling by dephosphorylating the phosphotryosine residues of insulin receptor kinase. This PTP was also reported to dephosphorylate epidermal growth factor receptor kinase, as well as JAK2 and TYK2 kinases, which implicated the role of this PTP in cell growth control, and cell response to interferon stimulation. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2013]

Protein Families: Druggable Genome, Phosphatase, Transmembrane

Protein Pathways: Adherens junction, Insulin signaling pathway

Product images:

