

## **Product datasheet for TP710264**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MCK10 (DDR1) (NM\_001954) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human discoidin domain receptor tyrosine kinase 1 (DDR1),

transcript variant 1, full length, with C-terminal DDK tag, expressed in sf9, 20ug

Species: Human

**Expression Host:** Sf9

**Expression cDNA Clone** 

or AA Sequence:

A DNA sequence from TrueORF clone, RC201224, encoding human full-length DDR1

Tag: C-DDK

**Predicted MW:** 97 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 50 mM Tris-HCl, 100 mM glycine, pH 8.0, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001945

Locus ID: 780

UniProt ID: <u>Q08345</u>, <u>A0A024RCQ1</u>

RefSeq Size: 3840

Cytogenetics: 6p21.33

RefSeq ORF: 2628

Synonyms: CAK; CD167; DDR; EDDR1; HGK2; MCK10; NEP; NTRK4; PTK3; PTK3A; RTK6; TRKE





**Summary:** 

Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. The protein encoded by this gene belongs to a subfamily of tyrosine kinase receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2011]

**Protein Families:** 

Druggable Genome, Protein Kinase, Transmembrane

## **Product images:**

