

#### OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 2

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TP710371

## CBX1 (NM\_006807) Human Recombinant Protein

## **Product data:**

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of human chromobox homolog 1 (CBX1), transcript variant 1, full length, with C-terminal DDK tag, expressed in sf9, 20ug
Species:	Human
Expression Host:	Sf9
Expression cDNA Clone or AA Sequence:	A DNA sequence from TrueORF clone, RC205672, encoding human full-length CBX1
Tag:	C-DDK
Predicted MW:	21.2 kDa
Concentration:	>0.05 $\mu$ g/ $\mu$ L as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	50 mM Tris-HCl, 100 mM glycine, pH 8.0, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP 006798</u>
Locus ID:	10951
UniProt ID:	<u>P83916</u>
RefSeq Size:	2443
Cytogenetics:	17q21.32
RefSeq ORF:	555
Synonyms:	CBX; HP1-BETA; HP1Hs-beta; HP1Hsbeta; M31; MOD1; p25beta




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** CBX1 (NM\_006807) Human Recombinant Protein – TP710371

Summary:

This gene encodes a highly conserved nonhistone protein, which is a member of the heterochromatin protein family . The protein is enriched in the heterochromatin and associated with centromeres. The protein has a single N-terminal chromodomain which can bind to histone proteins via methylated lysine residues, and a C-terminal chromo shadow-domain (CSD) which is responsible for the homodimerization and interaction with a number of chromatin-associated nonhistone proteins. The protein may play an important role in the epigenetic control of chromatin structure and gene expression. Several related pseudogenes are located on chromosomes 1, 3, and X. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Jul 2008]

## **Product images:**



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US