

Product datasheet for **TP720008M**

IL6 (NM_000600) Human Recombinant Protein

Product data:

| | |
|---------------------------------------|---|
| Product Type: | Recombinant Proteins |
| Description: | Recombinant protein of human interleukin 6 (interferon, beta 2) (IL6), the mature form (Pro29-Met212) with C-terminal His tag |
| Species: | Human |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | Val30-Met212 |
| Tag: | Tag Free |
| Predicted MW: | 20.9 kDa |
| Concentration: | lot specific |
| Purity: | >95% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl |
| Bioactivity: | ED50 is less than 0.1 ng/ml as determined by the dose-dependent stimulation of human TF-1 cells. Specific Activity of 5.0 x 10 ⁷ IU/mg. |
| Endotoxin: | < 0.1 EU per µg protein as determined by LAL test |
| Reconstitution Method: | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH ₂ O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |
| Storage: | Store at -80°C. |
| Stability: | Stable for at least 6 months from date of receipt under proper storage and handling conditions. |
| RefSeq: | NP_000591 |
| Locus ID: | 3569 |
| UniProt ID: | P05231 , Q75MH2 , B4DVM1 |
| Cytogenetics: | 7p15.3 |
| Synonyms: | BSF-2; BSF2; CDF; HGF; HSF; IFN-beta-2; IFNB2; IL-6 |



[View online »](#)

Summary:

This gene encodes a cytokine that functions in inflammation and the maturation of B cells. In addition, the encoded protein has been shown to be an endogenous pyrogen capable of inducing fever in people with autoimmune diseases or infections. The protein is primarily produced at sites of acute and chronic inflammation, where it is secreted into the serum and induces a transcriptional inflammatory response through interleukin 6 receptor, alpha. The functioning of this gene is implicated in a wide variety of inflammation-associated disease states, including susceptibility to diabetes mellitus and systemic juvenile rheumatoid arthritis. Elevated levels of the encoded protein have been found in virus infections, including COVID-19 (disease caused by SARS-CoV-2). [provided by RefSeq, Aug 2020]

Protein Families:

Druggable Genome, Secreted Protein

Protein Pathways:

Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing pathway, Graft-versus-host disease, Hematopoietic cell lineage, Hypertrophic cardiomyopathy (HCM), Jak-STAT signaling pathway, NOD-like receptor signaling pathway, Pathways in cancer, Prion diseases, Toll-like receptor signaling pathway

Product images: