

## **Product datasheet for TP720009L**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Parathyroid Hormone (PTH) (NM\_000315) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human parathyroid hormone (PTH)

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

Ser32-Gln115

Tag: Tag Free
Predicted MW: 9.4 kDa
Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 10mM HAc-NaAc, 150mM NaCl, 5% Mannitol,

pH 4.0.

**Bioactivity:** Specific Activity of 1.0 x 104 IU/mg as calculated by the UMR106 cell/cAMP method.

**Endotoxin:** < 0.1 EU per μg protein as determined by LAL test

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100  $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** NP 000306

 Locus ID:
 5741

 UniProt ID:
 P01270

 Cytogenetics:
 11p15.3

 Synonyms:
 FIH1; PTH1





**Summary:** 

This gene encodes a member of the parathyroid family of proteins. The encoded preproprotein is proteolytically processed to generate a protein that binds to the parathyroid hormone/parathyroid hormone-related peptide receptor and regulates blood calcium and phosphate levels. Excess production of the encoded protein, known as hyperparathyroidism, can result in hypercalcemia and kidney stones. On the other hand, defective processing of the encoded protein may lead to hypoparathyroidism, which can result in hypocalcemia and numbness. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2015]

**Protein Families:** 

Druggable Genome, Secreted Protein

## **Product images:**

