

## Product datasheet for TP720025L

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL1RA (IL1RN) (NM\_000577) Human Recombinant Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human interleukin 1 receptor antagonist (IL1RN), transcript variant 3

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

Expression CDNA Cione

Arg26-Glu177

or AA Sequence:

Tag: Tag Free
Predicted MW: 17.3 kDa
Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Bioactivity: ED50 is 0.5 ng/ml as determined by the dose-dependent inhibition of IL-1 stimulation of D10S

cells. Specific Activity of 2.0 x 106 IU/mg.

**Endotoxin:** < 0.1 EU per μg protein as determined by LAL test

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Storage:** Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeg: NP 000568

 Locus ID:
 3557

 UniProt ID:
 P18510

 Cytogenetics:
 2q14.1

Synonyms: DIRA; ICIL-1RA; IL-1ra; IL-1ra3; IL-1RN; IL1F3; IL1RA; IRAP; MVCD4





**Summary:** 

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This protein inhibits the activities of interleukin 1, alpha (IL1A) and interleukin 1, beta (IL1B), and modulates a variety of interleukin 1 related immune and inflammatory responses, particularly in the acute phase of infection and inflammation. This gene and five other closely related cytokine genes form a gene cluster spanning approximately 400 kb on chromosome 2. A polymorphism of this gene is reported to be associated with increased risk of osteoporotic fractures and gastric cancer. Several alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Aug 2020]

**Protein Families:** 

Druggable Genome, Secreted Protein

# **Product images:**

