

## **Product datasheet for TP720047**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CXCL14 (NM\_004887) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human chemokine (C-X-C motif) ligand 14 (CXCL14)

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

Ser35-Glu111

or AA Sequence:

Tag: Tag Free
Predicted MW: 9.4 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

**Bioactivity:** ED50 is 1.0-10.0 ng/ml as determined by the ability of Recombinant CXCL14 to induce

calcium flux of prostaglandin E2 treated THP1 human acute monocytic leukemia cells.

**Endotoxin:** < 0.1 EU per μg protein as determined by LAL test

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Storage:** Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 004878

**Locus ID:** 9547

 UniProt ID:
 O95715

 Cytogenetics:
 5q31.1

Synonyms: BMAC; BRAK; KEC; KS1; MIP-2g; MIP2G; NJAC; SCYB14



**Summary:** 

This antimicrobial gene belongs to the cytokine gene family which encode secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC (Cys-X-Cys) subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes but not for lymphocytes, dendritic cells, neutrophils or macrophages. It has been implicated that this cytokine is involved in the homeostasis of monocyte-derived macrophages rather than in inflammation. [provided by RefSeq, Sep 2014]

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

## **Product images:**

