

Product datasheet for **TP720053L**

CCL27 (NM_006664) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human chemokine (C-C motif) ligand 27 (CCL27)
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Phe25-Gly112
Tag:	Tag Free
Predicted MW:	10.1 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	< 0.1 EU per µg protein as determined by LAL test
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH ₂ O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	NP_006655
Locus ID:	10850
UniProt ID:	Q9Y4X3 , Q5VZ77
Cytogenetics:	9p13.3
Synonyms:	ALP; CTACK; CTAK; ESKINE; ILC; PESKY; SCYA27



[View online »](#)

Summary:

This gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene is chemotactic for skin-associated memory T lymphocytes. This cytokine may also play a role in mediating homing of lymphocytes to cutaneous sites. It specifically binds to chemokine receptor 10 (CCR10). Studies of a similar murine protein indicate that these protein-receptor interactions have a pivotal role in T cell-mediated skin inflammation. [provided by RefSeq, Sep 2014]

Protein Families:

Druggable Genome

Protein Pathways:

Chemokine signaling pathway, Cytokine-cytokine receptor interaction

Product images: