

Product datasheet for TP720083

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

beta Arrestin 1 (ARRB1) (NM 004041) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human arrestin, beta 1 (ARRB1), transcript variant 1

Species: Human E. coli **Expression Host:**

Expression cDNA Clone

Met1-Arg418

or AA Sequence:

C-His

Tag: Predicted MW: 48.1 kDa **Concentration:** lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: < 0.1 EU per µg protein as determined by LAL test

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

> lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Store at -80°C. Storage:

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

NP 004032 RefSeq:

Locus ID: 408

UniProt ID: P49407, B7Z1Q3

Cytogenetics: 11q13.4 Synonyms: ARB1; ARR1





Summary:

Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of G-protein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 1 is a cytosolic protein and acts as a cofactor in the beta-adrenergic receptor kinase (BARK) mediated desensitization of beta-adrenergic receptors. Besides the central nervous system, it is expressed at high levels in peripheral blood leukocytes, and thus the BARK/beta-arrestin system is believed to play a major role in regulating receptor-mediated immune functions. Alternatively spliced transcripts encoding different isoforms of arrestin beta 1 have been described. [provided by RefSeq, Jan 2011]

Protein Families: Druggable Genome

Protein Pathways: Chemokine signaling pathway, Endocytosis, MAPK signaling pathway

Product images:

