

## Product datasheet for **TP720084M**

### **BCL2A1 (NM\_001114735) Human Recombinant Protein**

#### **Product data:**

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human BCL2-related protein A1 (BCL2A1), transcript variant 2
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Met1-Ser152
Tag:	C-His
Predicted MW:	18.5 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	#N/A
Endotoxin:	< 0.1 EU per µg protein as determined by LAL test
Storage:	Store at -80°C.
Stability:	Stable for at least 3 months from date of receipt under proper storage and handling conditions.
RefSeq:	<u><a href="#">NP_001108207</a></u>
Locus ID:	597
UniProt ID:	<u><a href="#">Q16548</a></u>
Cytogenetics:	15q25.1
Synonyms:	ACC-1; ACC-2; ACC1; ACC2; BCL2L5; BFL1; GRS; HBPA1



[View online »](#)

**Summary:**

This gene encodes a member of the BCL-2 protein family. The proteins of this family form hetero- or homodimers and act as anti- and pro-apoptotic regulators that are involved in a wide variety of cellular activities such as embryonic development, homeostasis and tumorigenesis. The protein encoded by this gene is able to reduce the release of pro-apoptotic cytochrome c from mitochondria and block caspase activation. This gene is a direct transcription target of NF-kappa B in response to inflammatory mediators, and is up-regulated by different extracellular signals, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), CD40, phorbol ester and inflammatory cytokine TNF and IL-1, which suggests a cytoprotective function that is essential for lymphocyte activation as well as cell survival. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

**Protein Families:**

Druggable Genome

**Protein Pathways:**

Metabolic pathways

**Product images:**
