

Product datasheet for TP720085XL

Bcl x (BCL2L1) (NM_001191) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins Description: Recombinant protein of human BCL2-like 1 (BCL2L1), nuclear gene encoding mitochondrial protein, transcript variant 2 Species: Human **Expression Host:** E. coli **Expression cDNA Clone** Met1-Arg212 or AA Sequence: C-His Tag: Predicted MW: 24.8 kDa **Concentration:** lot specific **Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl Endotoxin: < 0.1 EU per µg protein as determined by LAL test Store at -80°C. Storage: Stable for at least 3 months from date of receipt under proper storage and handling Stability: conditions. RefSeq: NP 001182 Locus ID: 598 **UniProt ID:** Q07817 Cytogenetics: 20q11.21 Synonyms: Bcl-X; BCL-XL/S; BCL2L; BCLX; PPP1R52



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| ORIGENE BCI | x (BCL2L1) (NM_001191) Human Recombinant Protein – TP720085XL |
|--------------------|---|
| Summary: | The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform acts as an apoptotic inhibitor and the shorter isoform acts as an apoptotic activator. [provided by RefSeq, Dec 2015] |
| Protein Families: | Druggable Genome, ES Cell Differentiation/IPS, Transmembrane |
| Protein Pathways: | Amyotrophic lateral sclerosis (ALS), Apoptosis, Chronic myeloid leukemia, Jak-STAT signaling pathway, Pancreatic cancer, Pathways in cancer, Small cell lung cancer |

Product images:

~ 火



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US