

Product datasheet for TP720101XL

SDF1 (CXCL12) (NM_000609) Human Recombinant Protein

Product data:

| Product Type: | Recombinant Proteins |
|--|--|
| Description: | Recombinant protein of human chemokine (C-X-C motif) ligand 12 (stromal cell-derived factor 1) (CXCL12), transcript variant 2 |
| Species: | Human |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | Ser19-Met93 |
| Tag: | tag free |
| Predicted MW: | 8.8 kDa |
| Concentration: | lot specific |
| Purity: | >95% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | Lyophilized from a 0.2 um filtered solution of PBS, pH7.4. |
| Reconstitution Method: | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |
| Storage: | Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Stability: | Stable for at least 6 months from date of receipt under proper storage and handling conditions. |
| RefSeq: | <u>NP 000600</u> |
| Locus ID: | 6387 |
| UniProt ID: | <u>P48061</u> |
| RefSeq Size: | 3560 |
| Cytogenetics: | 10q11.21 |
| RefSeq ORF: | 279 |
| Synonyms: | IRH; PBSF; SCYB12; SDF1; TLSF; TPAR1 |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| | SDF1 (CXCL12) (NM_000609) Human Recombinant Protein – TP720101XL |
|-----------------|--|
| Summary: | This antimicrobial gene encodes a stromal cell-derived alpha chemokine member of the intercrine family. The encoded protein functions as the ligand for the G-protein coupled receptor, chemokine (C-X-C motif) receptor 4, and plays a role in many diverse cellular functions, including embryogenesis, immune surveillance, inflammation response, tissue homeostasis, and tumor growth and metastasis. Mutations in this gene are associated with resistance to human immunodeficiency virus type 1 infections. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2014] |
| Protein Familie | s: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein |
| Protein Pathwa | ys: Axon guidance, Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Leukocyte transendothelial migration |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US