

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP720118L

Fatty Acid Binding Protein 5 (FABP5) (NM_001444) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human fatty acid binding protein 5 (psoriasis-associated) (FABP5)
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Ala2-Glu135
Tag:	N-His
Predicted MW:	17.3 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 μ m filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	< 0.1 EU per µg protein as determined by LAL test
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<u>NP 001435</u>
Locus ID:	2171
UniProt ID:	<u>Q01469</u> , <u>E7DVW5</u>
Cytogenetics:	8q21.13
Synonyms:	E-FABP; EFABP; KFABP; PA-FABP; PAFABP




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Stty Acid Binding Protein 5 (FABP5) (NM_001444) Human Recombinant Protein – TP720118L

Summary:This gene encodes the fatty acid binding protein found in epidermal cells, and was first
identified as being upregulated in psoriasis tissue. Fatty acid binding proteins are a family of
small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other
hydrophobic ligands. FABPs may play roles in fatty acid uptake, transport, and metabolism.
Polymorphisms in this gene are associated with type 2 diabetes. The human genome
contains many pseudogenes similar to this locus.[provided by RefSeq, Feb 2011]

Protein Pathways: PPAR signaling pathway

Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US