

## Product datasheet for TP720154L

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Sumo 2 (SUMO2) (NM 001005849) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human SMT3 suppressor of mif two 3 homolog 2 (S. cerevisiae)

(SUMO2), transcript variant 2

Species: Human Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Met1-Gly93

Tag: N-His
Predicted MW: 13 kDa
Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

**Endotoxin:** < 0.1 EU per µg protein as determined by LAL test

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeg:** NP 001005849

 Locus ID:
 6613

 UniProt ID:
 P61956

 Cytogenetics:
 17q25.1

Synonyms: HSMT3; Smt3A; SMT3B; SMT3H2; SUMO3





**Summary:** 

This gene encodes a protein that is a member of the SUMO (small ubiquitin-like modifier) protein family. It functions in a manner similar to ubiquitin in that it is bound to target proteins as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as nuclear transport, transcriptional regulation, apoptosis, and protein stability. It is not active until the last two amino acids of the carboxy-terminus have been cleaved off. Numerous pseudogenes have been reported for this gene. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]

**Protein Families:** 

Druggable Genome

## **Product images:**

