

# **Product datasheet for TP720208**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# IMPA1 (NM 001144878) Human Recombinant Protein

#### **Product data:**

**Product Type: Recombinant Proteins** 

Description: Recombinant protein of human inositol(myo)-1(or 4)-monophosphatase 1 (IMPA1), transcript

variant 2

Species: Human **Expression Host:** E. coli

**Expression cDNA Clone** 

or AA Sequence:

Met1-Asp277

N-His Tag:

**Predicted MW:** 32.3 kDa Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Supplied as a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.25.

**Endotoxin:** < 0.1 EU per µg protein as determined by LAL test

Store at < -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles. Storage: Stability:

Stable for at least 3 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 001138350

Locus ID: 3612 **UniProt ID:** P29218 **Cytogenetics:** 8q21.13

Synonyms: IMP; IMPA; MRT59





**Summary:** 

This gene encodes an enzyme that dephosphorylates myo-inositol monophosphate to generate free myo-inositol, a precursor of phosphatidylinositol, and is therefore an important modulator of intracellular signal transduction via the production of the second messengers myoinositol 1,4,5-trisphosphate and diacylglycerol. This enzyme can also use myo-inositol-1,3-diphosphate, myo-inositol-1,4-diphosphate, scyllo-inositol-phosphate, glucose-1-phosphate, glucose-1-phosphate, fructose-1-phosphate, beta-glycerophosphate, and 2'-AMP as substrates. This enzyme shows magnesium-dependent phosphatase activity and is inhibited by therapeutic concentrations of lithium. Inhibition of inositol monophosphate hydroylosis and subsequent depletion of inositol for phosphatidylinositol synthesis may explain the anti-manic and anti-depressive effects of lithium administered to treat bipolar disorder. Alternative splicing results in multiple transcript variants encoding distinct isoforms. A pseudogene of this gene is also present on chromosome 8q21.13. [provided by RefSeq, Dec 2014]

**Protein Families:** Druggable Genome

Protein Pathways: Inositol phosphate metabolism, Metabolic pathways, Phosphatidylinositol signaling system

## **Product images:**

