

Product datasheet for TP720256M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PREI3 (MOB4) (NM_001100819) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human MOB1, Mps One Binder kinase activator-like 3 (yeast)

(MOBKL3), transcript variant 3

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Met1-Ala225

Tag: N-His

Predicted MW: 28.2 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM Tris-HCl, 150mM NaCl, 1mM DTT, pH 8.0.

Endotoxin: < 0.1 EU per µg protein as determined by LAL test

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeg: NP 001094289

 Locus ID:
 25843

 UniProt ID:
 Q9Y3A3

 Cytogenetics:
 2q33.1

Synonyms: 2C4D; CGI-95; MOB1; MOB3; MOBKL3; PHOCN; PREI3





Summary:

This gene was identified based on its similarity with the mouse counterpart. Studies of the mouse counterpart suggest that the expression of this gene may be regulated during oocyte maturation and preimplantation following zygotic gene activation. Alternatively spliced transcript variants encoding distinct isoforms have been observed. Naturally occurring readthrough transcription occurs between this locus and the neighboring locus HSPE1.[provided by RefSeq, Feb 2011]

Product images:

