

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP720261

PPIL1 (NM_016059) Human Recombinant Protein

Product data:


Description:Recombinant protein of human peptidylprolyl isomerase (cyclophilin)-like 1 (PPIL1)Species:HumanExpression Host:E. coliExpression cDNA Clome or AA Sequence:Met1-Gly166Tag:N-HisPredicted MW:20.4 kDaConcentration:Iot specificPurity:>95% as determined by SDS-PAGE and Coomasise blue stainingBuffer:Concentration:Storage:Concel Log program determined by LAL testStorage:Storage and Social	Product Type:	Recombinant Proteins
Expression Host:E. coliExpression CDNA CloueMet1-Gly166Frag:N-HisTag:0.4 kDaPredicted MW:0.4 kDaOncentration:Id specificPurity:>95% as determined by SDS-PAGE and Coomassie blue stainingBuffer:Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaClEndotoxin:<0.1 EU per µg protein as determined by LAL test	Description:	Recombinant protein of human peptidylprolyl isomerase (cyclophilin)-like 1 (PPIL1)
Presession cDNA ClowMet1-Gly166rag:N-HisTag:N-HisPredicted MW:20.4 kDaConcentration:lot specificPurity:>95% as determined by SDS-PAGE and Coomassie blue stainingBuffer:Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaClFhotoxin:< 0.1 EU per µg protein as determined by LAL test	Species:	Human
or AA Sequence:Tag:N-HisPredicted MW:20.4 kDaConcentration:lot specificPurity:>95% as determined by SDS-PAGE and Coomassie blue stainingBuffer:Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaClEndotoxin:< 0.1 EU per µg protein as determined by LAL test	Expression Host:	E. coli
Predicted MW:20.4 kDaPredicted MW:20.4 kDaConcentration:lot specificPurity:>95% as determined by SDS-PAGE and Coomassie blue stainingBuffer:Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaClEndotoxin:< 0.1 EU per µg protein as determined by LAL testStorage:Store at -80°C.Stability:Stable for at least 3 months from date of receipt under proper storage and handling conditions.RefSeq:NP 057143Locus ID:51645UniProt ID:Q9Y3C6, A0A024RCX8, A0A494C014Cytogenetics:Gp21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPlaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerase (PPlases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important ryclophilins are a highly conserved, ubiquitous family, members of which play an important ryclophilins are a highly conserved, ubiquitous family, members of which play an important ryclophilins are a highly conserved, ubiquitous family, members of which play an important might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	•	Met1-Gly166
Concentration:lot specificPurity:>95% as determined by SDS-PAGE and Coomassie blue stainingBuffer:Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaClEndotoxin:< 0.1 EU per µg protein as determined by LAL testStorage:Store at -80°C.Stability:Stable for at least 3 months from date of receipt under proper storage and handling conditions.RefSeq:NP 057143Locus ID:51645UniProt ID:09Y3C6, A0A024RCX8, A0A494C014Cytogenetics:621.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPlaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPlases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPlases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq. Jul 2008]	Tag:	N-His
Purity:>95% as determined by SDS-PAGE and Coomassie blue stainingBuffer:Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaClEndotoxin:< 0.1 EU per µg protein as determined by LAL testStorage:Store at -80°C.Stability:Stable for at least 3 months from date of receipt under proper storage and handling conditions.RefSeq:NP 057143Locus ID:51645UniProt ID:09Y3C6, A0A024RCX8, A0A494C014Cytogenetics:6p21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPIaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	Predicted MW:	20.4 kDa
Buffer:Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaClEndotoxin:< 0.1 EU per µg protein as determined by LAL testStorage:Store at -80°C.Stability:Stable for at least 3 months from date of receipt under proper storage and handling conditions.RefSeq:NP 057143Locus ID:51645UniProt ID:O9Y3C6, A0A024RCX8, A0A494C014Cytogenetics:6p21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPlaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPlases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPlases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq. Jul 2008]	Concentration:	lot specific
Endotoxin:< 0.1 EU per µg protein as determined by LAL test	Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Storage:Store at -80°C.Stability:Stable for at least 3 months from date of receipt under proper storage and handling conditions.RefSeq:NP 057143Locus ID:51645UniProt ID:Q9Y3C6, A0A024RCX8, A0A494C014Cytogenetics:6p21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPIaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	Buffer:	Provided lyophilized from a 0.2 μ m filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Stability:Stable for at least 3 months from date of receipt under proper storage and handling conditions.RefSeq:NP 057143Locus ID:51645UniProt ID:Q9Y3C6, A0A024RCX8, A0A494C014Cytogenetics:6p21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPIaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	Endotoxin:	< 0.1 EU per μ g protein as determined by LAL test
conditions.RefSeq:NP 057143Locus ID:51645UniProt ID:Q9Y3C6, A0A024RCX8, A0A494C014Cytogenetics:6p21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPIaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	Storage:	Store at -80°C.
Locus ID:51645UniProt ID:Q9Y3C6, A0A024RCX8, A0A494C014Cytogenetics:6p21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPIaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	Stability:	
UniProt ID:O9Y3C6, A0A024RCX8, A0A494C014Cytogenetics:6p21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPIaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	RefSeq:	<u>NP 057143</u>
Cytogenetics:6p21.2Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPIaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	Locus ID:	51645
Synonyms:CGI-124; CYPL1; hCyPX; PCH14; PPIaseSummary:This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	UniProt ID:	<u>Q9Y3C6, A0A024RCX8, A0A494C014</u>
Summary: This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	Cytogenetics:	6p21.2
cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]	Synonyms:	CGI-124; CYPL1; hCyPX; PCH14; PPlase
Protein Pathways: Spliceosome	Summary:	cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPlases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.
	Protein Pathways:	Spliceosome



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US