

Product datasheet for **TP720288M**

CRISP3 (NM_001190986) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cysteine-rich secretory protein 3 (CRISP3), transcript variant 2.
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	Asn21-Tyr245
Tag:	C-His
Predicted MW:	26.5 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.
Endotoxin:	< 0.1 EU per µg protein as determined by LAL test
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH ₂ O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	NP_001177915
Locus ID:	10321
UniProt ID:	P54108
Cytogenetics:	6p12.3
Synonyms:	Aeg2; CRISP-3; CRS3; dJ442L6.3; SGP28



[View online »](#)

Summary:

This gene encodes a member of the cysteine-rich secretory protein (CRISP) family within the CRISP, antigen 5 and pathogenesis-related 1 proteins superfamily. The encoded protein has an N-terminal CRISP, antigen 5 and pathogenesis-related 1 proteins domain, a hinge region, and a C-terminal ion channel regulator domain. This protein contains cysteine residues, located in both the N- and C-terminal domains, that form eight disulfide bonds, a distinguishing characteristic of this family. This gene is expressed in the male reproductive tract where it plays a role in sperm function and fertilization, and the female reproductive tract where it plays a role in endometrial receptivity for embryo implantation. This gene is upregulated in certain types of prostate cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2016]

Protein Families:

Secreted Protein

Product images: