

Product datasheet for TP720288XL

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CRISP3 (NM 001190986) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human cysteine-rich secretory protein 3 (CRISP3), transcript variant

2

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Asn21-Tyr245

Tag: C-His

Predicted MW: 26.5 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

Endotoxin: < 0.1 EU per µg protein as determined by LAL test

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeg: NP 001177915

 Locus ID:
 10321

 UniProt ID:
 P54108

 Cytogenetics:
 6p12.3

Synonyms: Aeg2; CRISP-3; CRS3; dJ442L6.3; SGP28





Summary:

This gene encodes a member of the cysteine-rich secretory protein (CRISP) family within the CRISP, antigen 5 and pathogenesis-related 1 proteins superfamily. The encoded protein has an N-terminal CRISP, antigen 5 and pathogenesis-related 1 proteins domain, a hinge region, and a C-terminal ion channel regulator domain. This protein contains cysteine residues, located in both the N- and C-terminal domains, that form eight disulfide bonds, a distinguishing characteristic of this family. This gene is expressed in the male reproductive tract where it plays a role in sperm function and fertilization, and the female reproductive tract where it plays a role in endometrial receptivity for embryo implantation. This gene is upregulated in certain types of prostate cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2016]

Protein Families:

Secreted Protein

Product images:

