

Product datasheet for TP720294L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cystatin SA (CST2) (NM 001322) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human cystatin SA (CST2)

Species: Human Expression Host: HEK293

Expression cDNA Clone

Trp21-Ala141

or AA Sequence:

Tag: C-His

Predicted MW: 15.4 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.

Endotoxin: < 0.1 EU per μg protein as determined by LAL test

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 001313

 Locus ID:
 1470

 UniProt ID:
 P09228

 Cytogenetics:
 20p11.21





Summary:

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a secreted thiol protease inhibitor found at high levels in saliva, tears and seminal plasma. [provided by RefSeq, Jul 2008]

Protein Families: Secreted Protein

Product images:

