

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TP720303

FLRT2 (NM_013231) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human fibronectin leucine rich transmembrane protein 2 (FLRT2)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	Cys36-Ser539
Tag:	C-His
Predicted MW:	57.3 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Endotoxin:	< 0.1 EU per μ g protein as determined by LAL test
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<u>NP 037363</u>
Locus ID:	23768
UniProt ID:	<u>O43155</u>
Cytogenetics:	14q31.3



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE FLRT2 (NM_013231) Human Recombinant Protein – TP720303

Summary:This gene encodes a member of the fibronectin leucine rich transmembrane (FLRT) family of
cell adhesion molecules, which regulate early embryonic vascular and neural development.
The encoded type I transmembrane protein has an extracellular region consisting of an N-
terminal leucine-rich repeat domain and a type 3 fibronectin domain, followed by a
transmembrane domain and a short C-terminal cytoplasmic tail domain. It functions as both
a homophilic cell adhesion molecule and a heterophilic chemorepellent through its
interaction with members of the uncoordinated-5 receptor family. Proteolytic removal of the
extracellular region controls the migration of neurons in the developing cortex. Alternative
splicing results in multiple transcript variants. [provided by RefSeq, Sep 2016]

Protein Families:

Druggable Genome, Transmembrane

Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US