

## Product datasheet for **TP720394XL**

### CD89 (FCAR) (NM\_002000) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human Fc fragment of IgA, receptor for (FCAR), transcript variant 1
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	Gln22-Asn227
Tag:	C-His
Predicted MW:	24.5 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	< 0.1 EU per µg protein as determined by LAL test
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH <sub>2</sub> O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<a href="#">NP_001991</a>
Locus ID:	2204
UniProt ID:	<a href="#">P24071</a> , <a href="#">Q92588</a>
Cytogenetics:	19q13.42
Synonyms:	CD89; CTB-61M7.2; FcalphaRI



[View online »](#)

**Summary:**

This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]

**Protein Families:**

Transmembrane

**Product images:**