

Product datasheet for TP720395XL

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GDNF Receptor alpha 1 (GFRA1) (NM 001145453) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human GDNF family receptor alpha 1 (GFRA1), transcript variant 3

Species: Human **HEK293 Expression Host:**

Expression cDNA Clone

or AA Sequence:

C-His Tag:

Predicted MW: 46.3 kDa **Concentration:** lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB,150mM NaCl,pH7.4.

Endotoxin: < 0.1 EU per µg protein as determined by LAL test

Asp25-Lys429

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

> lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 Storage:

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stable for at least 6 months from date of receipt under proper storage and handling Stability:

conditions.

RefSeq: NP 001138925

Locus ID: 2674 **UniProt ID:** P56159 **Cytogenetics:** 10q25.3

GDNFR; GDNFRA; GFR-ALPHA-1; GFRalpha-1; RET1L; RETL1; TRNR1 Synonyms:





Summary:

This gene encodes a member of the glial cell line-derived neurotrophic factor receptor (GDNFR) family of proteins. The encoded preproprotein is proteolytically processed to generate the mature receptor. Glial cell line-derived neurotrophic factor (GDNF) and neurturin (NTN) are two structurally related, potent neurotrophic factors that play key roles in the control of neuron survival and differentiation. This receptor is a glycosylphosphatidylinositol (GPI)-linked cell surface receptor for both GDNF and NTN, and mediates activation of the RET tyrosine kinase receptor. This gene is a candidate gene for Hirschsprung disease. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed. [provided by RefSeq, Jan 2016]

Protein Families:

Druggable Genome

Product images:

