

Product datasheet for TP720398XL

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ICOS Ligand (ICOSLG) (NM_015259) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human inducible T-cell co-stimulator ligand (ICOSLG)

Species: Human
Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Asp19-Ser258

Tag: C-His
Predicted MW: 27.7 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: < 0.1 EU per μg protein as determined by LAL test

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 056074

Locus ID: 23308

UniProt ID: <u>075144</u>, <u>A0N0L8</u>

Cytogenetics: 21q22.3

Synonyms: B7-H2; B7h2; B7RP-1; B7RP1; CD275; GL50; ICOS-L; ICOSL; LICOS







Summary: Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-

cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation

into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-

stimulating memory T-cell function (By similarity).[UniProtKB/Swiss-Prot Function]

Protein Families: Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs)

Product images:

