

## Product datasheet for **TP720404**

### **LILRB2 (NM\_001080978) Human Recombinant Protein**

#### **Product data:**

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Recombinant protein of human leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 2 (LILRB2), transcript variant 2
<b>Species:</b>	Human
<b>Expression Host:</b>	HEK293
<b>Expression cDNA Clone or AA Sequence:</b>	Gln22-His458
<b>Tag:</b>	C-His
<b>Predicted MW:</b>	48.6 kDa
<b>Concentration:</b>	lot specific
<b>Purity:</b>	>95% as determined by SDS-PAGE and Coomassie blue staining
<b>Buffer:</b>	Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
<b>Endotoxin:</b>	< 0.1 EU per µg protein as determined by LAL test
<b>Reconstitution Method:</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH <sub>2</sub> O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
<b>Storage:</b>	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Stability:</b>	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
<b>RefSeq:</b>	<a href="#">NP_001074447</a>
<b>Locus ID:</b>	10288
<b>UniProt ID:</b>	<a href="#">A2IXV5</a>
<b>Cytogenetics:</b>	19q13.42
<b>Synonyms:</b>	CD85D; ILT-4; ILT4; LIR-2; LIR2; MIR-10; MIR10


[View online »](#)

**Summary:**

This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

**Protein Families:**

Transmembrane

**Product images:**
