

Product datasheet for TP720428XL

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CCL3L3 (NM_001001437) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human chemokine (C-C motif) ligand 3-like 3 (CCL3L3)

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Ala24-Ala93

Tage

Tag: C-His
Predicted MW: 8.8 kDa
Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB,150mM NaCl,pH7.4.

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 001001437

 Locus ID:
 414062

 UniProt ID:
 P16619

 RefSeq Size:
 780

 Cytogenetics:
 17q12

RefSeq ORF: 279

Synonyms: 464.2; D17S1718; G0S19-2; LD78; LD78BETA; SCYA3L; SCYA3L1





CCL3L3 (NM_001001437) Human Recombinant Protein - TP720428XL

Summary: This gene is one of several cytokine genes that are clustered on the q-arm of chromosome

17. Cytokines are a family of secreted proteins that function in inflammatory and

immunoregulatory processes. The protein encoded by this gene binds to several chemokine receptors, including chemokine binding protein 2 and chemokine (C-C motif) receptor 5 (CCR5). CCR5 is a co-receptor for HIV, and binding of this protein to CCR5 inhibits HIV entry. The copy number of this gene varies among individuals, where most individuals have one to six copies, and a minority of individuals have zero or more than six copies. There are conflicting reports about copy number variation of this gene and its correlation to disease

susceptibility.[provided by RefSeq, Apr 2014]

Protein Families: Druggable Genome

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction