

Product datasheet for TP720438L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PSP (REG1A) (NM 002909) Human Recombinant Protein

Gln23-Asn166

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human regenerating islet-derived 1 alpha (REG1A)

Species: Human **HEK293 Expression Host:**

Expression cDNA Clone

or AA Sequence:

C-His Tag:

Predicted MW: 17.3 kDa **Concentration:** lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Endotoxin: < 0.1 EU per µg protein as determined by LAL test

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

> lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 Storage:

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stable for at least 6 months from date of receipt under proper storage and handling Stability:

conditions.

RefSeq: NP 002900

Locus ID: 5967

UniProt ID: P05451, A8K7G6

Cytogenetics: 2p12

ICRF; P19; PSP; PSPS; PSPS1; PTP; REG Synonyms:





Summary:

This gene is a type I subclass member of the Reg gene family. The Reg gene family is a multigene family grouped into four subclasses, types I, II, III and IV, based on the primary structures of the encoded proteins. This gene encodes a protein that is secreted by the exocrine pancreas. It is associated with islet cell regeneration and diabetogenesis and may be involved in pancreatic lithogenesis. Reg family members REG1B, REGL, PAP and this gene are tandemly clustered on chromosome 2p12 and may have arisen from the same ancestral gene by gene duplication. [provided by RefSeq, Jul 2008]

Product images:

