

Product datasheet for TP720505L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LDL Receptor (LDLR) (NM 000527) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human low density lipoprotein receptor (LDLR), Ala22 - Arg788

Species: Human **HEK293 Expression Host:**

Expression cDNA Clone

or AA Sequence:

C-His Tag: Predicted MW: 86 kDa **Concentration:** lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: < 0.1 EU per µg protein as determined by LAL test

Ala22-Arg788

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

> lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Store at -80°C. Storage:

Stability: Stable for at least 3 months from date of receipt under proper storage and handling

conditions.

NP 000518 RefSeq:

Locus ID: 3949

UniProt ID: P01130, A0A024R7D5

Cytogenetics: 19p13.2

Synonyms: FH; FHC; FHCL1; LDLCQ2





Summary:

The low density lipoprotein receptor (LDLR) gene family consists of cell surface proteins involved in receptor-mediated endocytosis of specific ligands. Low density lipoprotein (LDL) is normally bound at the cell membrane and taken into the cell ending up in lysosomes where the protein is degraded and the cholesterol is made available for repression of microsomal enzyme 3-hydroxy-3-methylglutaryl coenzyme A (HMG CoA) reductase, the rate-limiting step in cholesterol synthesis. At the same time, a reciprocal stimulation of cholesterol ester synthesis takes place. Mutations in this gene cause the autosomal dominant disorder, familial hypercholesterolemia. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Sep 2010]

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Endocytosis

Product images:

