

## Product datasheet for TP720524M

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## WBP2 (NM 012478) Human Recombinant Protein

**Product data:** 

**Product Type: Recombinant Proteins** 

Description: Recombinant protein of human WW domain binding protein 2 (WBP2)

Species: Human E. coli **Expression Host:** 

**Expression cDNA Clone** 

Met1-Ala100

or AA Sequence:

N-His Tag:

Predicted MW: 13.4 kDa **Concentration:** lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Lyophilized from a 0.2 um filtered solution of 20mMTris-HCl,1mMDTT,5%Trehalose,pH8.0.

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

> lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 036610

Locus ID: 23558

**UniProt ID:** Q969T9, A0A024R8L1

RefSeq Size: 1944 Cytogenetics: 17q25.1

RefSeq ORF: 783

Synonyms: DFNB107; GRAMD6; WBP-2







**Summary:** 

The globular WW domain is composed of 38 to 40 semiconserved amino acids shared by proteins of diverse functions including structural, regulatory, and signaling proteins. The domain is involved in mediating protein-protein interactions through the binding of polyproline ligands. This gene encodes a WW domain binding protein that is a transcriptional coactivator of estrogen receptor alpha and progesterone receptor. Defects in this gene have been associated with hearing impairment. [provided by RefSeq, Jan 2017]