

## **Product datasheet for TP720535**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## GRB2 (NM\_002086) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human growth factor receptor-bound protein 2 (GRB2), transcript

variant 1

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Met1-Val217

Tag: C-His

Predicted MW: 26.3 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.

**Endotoxin:** < 0.1 EU per µg protein as determined by LAL test

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeg: NP 002077

 Locus ID:
 2885

 UniProt ID:
 P62993

 Cytogenetics:
 17q25.1

Synonyms: ASH; EGFRBP-GRB2; Grb3-3; MST084; MSTP084; NCKAP2





**Summary:** The protein encoded by this gene binds the epidermal growth factor receptor and contains

one SH2 domain and two SH3 domains. Its two SH3 domains direct complex formation with proline-rich regions of other proteins, and its SH2 domain binds tyrosine phosphorylated sequences. This gene is similar to the Sem5 gene of C.elegans, which is involved in the signal transduction pathway. Two alternatively spliced transcript variants encoding different

isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome

**Protein Pathways:** Acute myeloid leukemia, B cell receptor signaling pathway, Chemokine signaling pathway,

Chronic myeloid leukemia, Colorectal cancer, Dorso-ventral axis formation, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Focal adhesion, Gap junction, Glioma, GnRH signaling pathway, Insulin signaling pathway, Jak-STAT signaling pathway, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Non-small cell lung cancer, Pathways in cancer, Prostate cancer, Renal cell

carcinoma, T cell receptor signaling pathway

## **Product images:**

