

Product datasheet for **TP720543XL**

Mortality Factor 4 like 2 (MORF4L2) (NM_001142418) Human Recombinant Protein

Product data:

| | |
|---------------------------------------|---|
| Product Type: | Recombinant Proteins |
| Description: | Recombinant protein of human mortality factor 4 like 2 (MORF4L2), transcript variant 1 |
| Species: | Human |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | Met1-Leu288 |
| Tag: | C-His |
| Predicted MW: | 33.4 kDa |
| Concentration: | lot specific |
| Purity: | >95% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl |
| Endotoxin: | < 0.1 EU per µg protein as determined by LAL test |
| Reconstitution Method: | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH ₂ O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |
| Storage: | Store at -80°C. |
| Stability: | Stable for at least 6 months from date of receipt under proper storage and handling conditions. |
| RefSeq: | NP_001135890 |
| Locus ID: | 9643 |
| UniProt ID: | B3KP92 , Q15014 |
| Cytogenetics: | Xq22.2 |
| Synonyms: | MORFL2; MRGX |



[View online »](#)

Summary:

Component of the NuA4 histone acetyltransferase complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histone H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the MSIN3A complex which acts to repress transcription by deacetylation of nucleosomal histones. [UniProtKB/Swiss-Prot Function]

Protein Families:

Transcription Factors

Product images: