

Product datasheet for **TP720563M**

BCAS2 (NM_005872) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human breast carcinoma amplified sequence 2 (BCAS2)
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Ala2-Phe225
Tag:	N-T7&C-His
Predicted MW:	28.6 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	< 0.1 EU per µg protein as determined by LAL test
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH ₂ O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	NP_005863
Locus ID:	10286
UniProt ID:	O75934 , B2R7W3
Cytogenetics:	1p13.2
Synonyms:	DAM1; Snt309; SPF27



[View online »](#)

Summary:

Required for pre-mRNA splicing as component of the activated spliceosome (PubMed:28502770, PubMed:28076346, PubMed:29360106, PubMed:29301961, PubMed:30705154). Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing. May have a scaffolding role in the spliceosome assembly as it contacts all other components of the core complex. The PRP19-CDC5L complex may also play a role in the response to DNA damage (DDR). [UniProtKB/Swiss-Prot Function]

Protein Pathways:

Spliceosome

Product images: