

Product datasheet for TP720609XL

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Oncostatin M (OSM) (NM_020530) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human oncostatin M (OSM)

Species: Human
Expression Host: E. coli

Expression cDNA Clone

Expression CDNA Cion

Ala26-Arg221

or AA Sequence:

Tag: N-His

Predicted MW: 24.4 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 065391

 Locus ID:
 5008

 UniProt ID:
 P13725

 RefSeq Size:
 1869

 Cytogenetics:
 22q12.2

RefSeq ORF: 756





Oncostatin M (OSM) (NM_020530) Human Recombinant Protein - TP720609XL

Summary:

This gene encodes a member of the leukemia inhibitory factor/oncostatin-M (LIF/OSM) family of proteins. The encoded preproprotein is proteolytically processed to generate the mature protein. This protein is a secreted cytokine and growth regulator that inhibits the proliferation of a number of tumor cell lines. This protein also regulates the production of other cytokines, including interleukin 6, granulocyte-colony stimulating factor and granulocyte-macrophage colony stimulating factor in endothelial cells. This gene and the related gene, leukemia inhibitory factor, also present on chromosome 22, may have resulted from the duplication of a common ancestral gene. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Jan 2016]

Protein Families:

 $\label{lem:condition} \textit{Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein, Stem cell relevant signaling -- \\$

DSL/Notch pathway, Stem cell relevant signaling - JAK/STAT signaling pathway

Protein Pathways:

Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway