

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TP720614

## LTA (NM\_000595) Human Recombinant Protein

## **Product data:**

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human lymphotoxin alpha (TNF superfamily, member 1) (LTA), transcript variant 2
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	Leu35-Leu205
Tag:	Tag Free
Predicted MW:	18.8 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 $\mu$ m filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Reconstitution Method: Storage:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less
	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling
Storage: Stability:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions.
Storage: Stability: RefSeq:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 000586</u>
Storage: Stability: RefSeq: Locus ID:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 000586</u> 4049
Storage: Stability: RefSeq: Locus ID: UniProt ID:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 000586</u> 4049 <u>P01374, Q5STV3</u>
Storage: Stability: RefSeq: Locus ID: UniProt ID: RefSeq Size:	lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling conditions. <u>NP 000586</u> 4049 <u>P01374, Q5STV3</u> 1423



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	LTA (NM_000595) Human Recombinant Protein – TP720614
Summary:	The encoded protein, a member of the tumor necrosis factor family, is a cytokine produced by lymphocytes. The protein is highly inducible, secreted, and forms heterotrimers with lymphotoxin-beta which anchor lymphotoxin-alpha to the cell surface. This protein also mediates a large variety of inflammatory, immunostimulatory, and antiviral responses, is involved in the formation of secondary lymphoid organs during development and plays a role in apoptosis. Genetic variations in this gene are associated with susceptibility to leprosy type 4, myocardial infarction, non-Hodgkin's lymphoma, and psoriatic arthritis. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, Jul 2012]
Protein Families:	Druggable Genome, Secreted Protein, Transmembrane
Protein Pathway	<b>s:</b> Antigen processing and presentation, Cytokine-cytokine receptor interaction, Type I diabetes mellitus

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US