

## Product datasheet for **TP720621XL**

### **NDRG1 (NM\_006096) Human Recombinant Protein**

#### **Product data:**

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Purified recombinant protein of Human N-myc downstream regulated 1 (NDRG1), transcript variant 2
<b>Species:</b>	Human
<b>Expression Host:</b>	E. coli
<b>Expression cDNA Clone or AA Sequence:</b>	Met1-Cys394
<b>Tag:</b>	N-His
<b>Predicted MW:</b>	45 kDa
<b>Concentration:</b>	lot specific
<b>Purity:</b>	>95% as determined by SDS-PAGE and Coomassie blue staining
<b>Buffer:</b>	Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
<b>Endotoxin:</b>	Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)
<b>Reconstitution Method:</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH <sub>2</sub> O. It is not recommended to reconstitute a concentration less than 100 μg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
<b>Storage:</b>	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Stability:</b>	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
<b>RefSeq:</b>	<a href="#">NP_006087</a>
<b>Locus ID:</b>	10397
<b>UniProt ID:</b>	<a href="#">Q92597</a>
<b>RefSeq Size:</b>	3123
<b>Cytogenetics:</b>	8q24.22
<b>RefSeq ORF:</b>	1182



[View online »](#)

**Synonyms:** CAP43; CMT4D; DRG-1; DRG1; GC4; HMSNL; NDR1; NMSL; PROXY1; RIT42; RTP; TARG1; TDD5

**Summary:** This gene is a member of the N-myc downregulated gene family which belongs to the alpha/beta hydrolase superfamily. The protein encoded by this gene is a cytoplasmic protein involved in stress responses, hormone responses, cell growth, and differentiation. The encoded protein is necessary for p53-mediated caspase activation and apoptosis. Mutations in this gene are a cause of Charcot-Marie-Tooth disease type 4D, and expression of this gene may be a prognostic indicator for several types of cancer. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, May 2012]