

Product datasheet for **TP720631**

SLAMF7 (NM_021181) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human SLAM family member 7 (SLAMF7)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	Ser23-Met226
Tag:	C-His
Predicted MW:	23.3 kDa
Concentration:	N/A
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 μ m filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4
Endotoxin:	Endotoxin level is < 0.1 ng/ μ g of protein (< 1 EU/ μ g)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH ₂ O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	NP_067004
Locus ID:	57823
UniProt ID:	Q9NQ25
RefSeq Size:	2672
Cytogenetics:	1q23.3
RefSeq ORF:	1005
Synonyms:	19A; CD319; CRACC; CS1



[View online »](#)

Summary:

Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2. Isoform 1 mediates NK cell activation through a SH2D1A-independent extracellular signal-regulated ERK-mediated pathway (PubMed:11698418). Positively regulates NK cell functions by a mechanism dependent on phosphorylated SH2D1B. Downstream signaling implicates PLCG1, PLCG2 and PI3K (PubMed:16339536). In addition to heterotypic NK cells-target cells interactions also homotypic interactions between NK cells may contribute to activation. However, in the absence of SH2D1B, inhibits NK cell function. Acts also inhibitory in T-cells (By similarity). May play a role in lymphocyte adhesion (PubMed:11802771). In LPS-activated monocytes negatively regulates production of proinflammatory cytokines (PubMed:23695528).[UniProtKB/Swiss-Prot Function]

Protein Families:

Druggable Genome, Transmembrane