

Product datasheet for TP720632L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD32A (FCGR2A) (NM_021642) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human Fc fragment of IgG, low affinity Ila, receptor (CD32)

(FCGR2A), transcript variant 2

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Ala36-Ile218

Tag: C-His

Predicted MW: 21.61 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 067674

 Locus ID:
 2212

 UniProt ID:
 P12318

 RefSeq Size:
 2411

 Cytogenetics:
 1q23.3

 RefSeq ORF:
 948

Synonyms: CD32; CD32A; CDw32; FCG2; FcGR2; FCGR2A1; IGFR2





CD32A (FCGR2A) (NM_021642) Human Recombinant Protein - TP720632L

Summary: This gene encodes one member of a family of immunoglobulin Fc receptor genes found on

the surface of many immune response cells. The protein encoded by this gene is a cell surface receptor found on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes. Alternative

splicing results in multiple transcript variants. [provided by RefSeq, Oct 2008]

Protein Families: ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Fc gamma R-mediated phagocytosis, Systemic lupus erythematosus