

Product datasheet for TP720637

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DR6 (TNFRSF21) (NM 014452) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human tumor necrosis factor receptor superfamily, member

21 (TNFRSF21)

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Gln42-Leu350

Tag: C-His

Predicted MW: 34.62 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 μ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: <u>NP 055267</u>

Locus ID: 27242

UniProt ID: <u>075509</u>, <u>A0A024RD71</u>

RefSeq Size: 3646 Cytogenetics: 6p12.3 RefSeq ORF: 1965

Synonyms: BM-018; CD358; DR6





DR6 (TNFRSF21) (NM_014452) Human Recombinant Protein - TP720637

Summary: This gene encodes a member of the tumor necrosis factor receptor superfamily. The encoded

protein activates nuclear factor kappa-B and mitogen-activated protein kinase 8 (also called c-Jun N-terminal kinase 1), and induces cell apoptosis. Through its death domain, the encoded receptor interacts with tumor necrosis factor receptor type 1-associated death domain (TRADD) protein, which is known to mediate signal transduction of tumor necrosis factor

receptors. Knockout studies in mice suggest that this gene plays a role in T-helper cell activation, and may be involved in inflammation and immune regulation. [provided by

RefSeq, Jul 2013]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction