

## Product datasheet for **TP720640L**

### PROCR (NM\_006404) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human protein C receptor, endothelial (PROCR)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	Ser18-Ser210
Tag:	C-His
Predicted MW:	23.06 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Provided lyophilized from a 0.2 $\mu$ m filtered solution of 20 mM Tris-HCl, 150 mM NaCl
Endotoxin:	Endotoxin level is < 0.1 ng/ $\mu$ g of protein (< 1 EU/ $\mu$ g)
Reconstitution Method:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH <sub>2</sub> O. It is not recommended to reconstitute a concentration less than 100 $\mu$ g/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	<a href="#">NP_006395</a>
Locus ID:	10544
UniProt ID:	<a href="#">Q9UNN8</a>
RefSeq Size:	1506
Cytogenetics:	20q11.22
RefSeq ORF:	714
Synonyms:	CCCA; CCD41; EPCR



[View online »](#)

**Summary:**

The protein encoded by this gene is a receptor for activated protein C, a serine protease activated by and involved in the blood coagulation pathway. The encoded protein is an N-glycosylated type I membrane protein that enhances the activation of protein C. Mutations in this gene have been associated with venous thromboembolism and myocardial infarction, as well as with late fetal loss during pregnancy. The encoded protein may also play a role in malarial infection and has been associated with cancer. [provided by RefSeq, Jul 2013]

**Protein Families:**

Druggable Genome, ES Cell Differentiation/IPS, Transmembrane