

## **Product datasheet for TP720651**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SIRP alpha (SIRPA) (NM\_080792) Human Recombinant Protein

## **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human signal-regulatory protein alpha (SIRPA), transcript

variant 3

Species: Human Expression Host: HEK293

**Expression cDNA Clone** 

or AA Sequence:

Glu31-Arg370

Tag: C-His

Predicted MW: 38.31 kDa

Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Lyophilized from a 0.2 um filtered solution of PBS,PH7.4.

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** NP 542970

 Locus ID:
 140885

 UniProt ID:
 P78324

 RefSeq Size:
 3868

 Cytogenetics:
 20p13

RefSeq ORF: 1509

Synonyms: BIT; CD172A; MFR; MYD-1; P84; PTPNS1; SHPS1; SIRP



ORIGENE

**Summary:** 

The protein encoded by this gene is a member of the signal-regulatory-protein (SIRP) family, and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transmembrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. This protein can be phosphorylated by tyrosine kinases. The phospho-tyrosine residues of this PTP have been shown to recruit SH2 domain containing tyrosine phosphatases (PTP), and serve as substrates of PTPs. This protein was found to participate in signal transduction mediated by various growth factor receptors. CD47 has been demonstrated to be a ligand for this receptor protein. This gene and its product share very high similarity with several other members of the SIRP family. These related genes are located in close proximity to each other on chromosome 20p13. Multiple alternatively spliced transcript variants have been determined for this gene. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome, Phosphatase, Transmembrane