

## Product datasheet for TP720652L

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## LY108 (SLAMF6) (NM 052931) Human Recombinant Protein

**Product data:** 

**Product Type: Recombinant Proteins** 

Description: Purified recombinant protein of Human SLAM family member 6 (SLAMF6), transcript variant 2

Species: Human **HEK293 Expression Host:** 

**Expression cDNA Clone** 

or AA Sequence:

Leu28-Lys225

C-His Tag:

Predicted MW: 23.98 kDa **Concentration:** lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

**Endotoxin:** Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)

**Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

> lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Store at -80°C. Storage:

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

NP 443163 RefSeq: Locus ID: 114836

**UniProt ID:** Q96DU3

RefSeg Size: 2776

Cytogenetics: 1q23.2-q23.3

RefSeq ORF: 993

Synonyms: CD352; KALI; KALIb; Ly108; NTB-A; NTBA; SF2000





## LY108 (SLAMF6) (NM\_052931) Human Recombinant Protein - TP720652L

**Summary:** 

The protein encoded by this gene is a type I transmembrane protein, belonging to the CD2 subfamily of the immunoglobulin superfamily. This encoded protein is expressed on Natural killer (NK), T, and B lymphocytes. It undergoes tyrosine phosphorylation and associates with the Src homology 2 domain-containing protein (SH2D1A) as well as with SH2 domain-containing phosphatases (SHPs). It functions as a coreceptor in the process of NK cell activation. It can also mediate inhibitory signals in NK cells from X-linked lymphoproliferative patients. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, May 2010]

**Protein Families:** 

Druggable Genome, Transmembrane