

Product datasheet for TP720691L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD66b (CEACAM8) (NM_001816) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human carcinoembryonic antigen-related cell adhesion

molecule 8 (CEACAM8)

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Gln35-His141

Tag: C-His

Predicted MW: 13.03 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 001807

Locus ID: 1088

UniProt ID: <u>P31997</u>, <u>Q0Z7S6</u>, <u>B4DLI3</u>

RefSeq Size: 2333 Cytogenetics: 19q13.2

RefSeg ORF: 1047





CD66b (CEACAM8) (NM_001816) Human Recombinant Protein - TP720691L

Synonyms: CD66b; CD67; CGM6; NCA-95

Summary: Cell surface glycoprotein that plays a role in cell adhesion in a calcium-independent manner

(PubMed:8776764, PubMed:2022629, PubMed:11590190). Mediates heterophilic cell adhesion with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM6 (PubMed:8776764, PubMed:2022629, PubMed:11590190). Heterophilic interaction with CEACAM8 occurs in activated neutrophils (PubMed:8776764).[UniProtKB/Swiss-Prot

Function]