

Product datasheet for TP720755L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LILRA2 (NM_006866) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human leukocyte immunoglobulin-like receptor, subfamily A

(with TM domain), member 2 (LILRA2), transcript variant 2

Species: Human Expression Host: HEK293

Expression cDNA Clone

or AA Sequence:

Gly24-Ser420

Tag: C-His

Predicted MW: 44.8 kDa **Concentration:** lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 um filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Endotoxin: Endotoxin level is < 0.1 ng/μg of protein (< 1 EU/μg)

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3

weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 006857

 Locus ID:
 11027

 UniProt ID:
 Q8N149

 RefSeq Size:
 1696

Cytogenetics: 19q13.42

RefSeq ORF: 1398





LILRA2 (NM_006866) Human Recombinant Protein - TP720755L

Synonyms: CD85H; ILT1; LIR-7; LIR7

Summary: This gene encodes a member of a family of immunoreceptors that are expressed

predominantly on monocytes and B cells, and at lower levels on dendritic cells and natural killer cells. The encoded protein is an activating receptor that inhibits dendritic cell

differentiation and antigen presentation and suppresses innate immune response. Alternatively spliced transcript variants encoding different isoforms have been found. This gene is located in a cluster of related genes on chromosome 19 and there is a pseudogene

for this gene on chromosome 3. [provided by RefSeq, Mar 2014]