

# Product datasheet for TP720848XL

## BLNK (NM\_013314) Human Recombinant Protein

### **Product data:**

#### **Product Type: Recombinant Proteins Description:** Purified recombinant protein of Human B-cell linker (BLNK), transcript variant 1 Species: Human E. coli **Expression Host:** Met1-Ser456 **Expression cDNA Clone** or AA Sequence: C-His Tag: Predicted MW: 51.5 kDa **Concentration:** lot specific **Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl Endotoxin: Endotoxin level is < 0.1 ng/ $\mu$ g of protein (< 1 EU/ $\mu$ g) **Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Store at -80°C. Storage: Stability: Stable for at least 6 months from date of receipt under proper storage and handling conditions. NP 037446 RefSeq: Locus ID: 29760 UniProt ID: Q8WV28 **RefSeq Size:** 1829 Cytogenetics: 10q24.1 **RefSeq ORF:** 1368 Synonyms: AGM4; BASH; bca; BLNK-S; LY57; SLP-65; SLP65



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	BLNK (NM_013314) Human Recombinant Protein – TP720848XL
Summary:	This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, May 2012]
Protein Families Protein Pathway	
-	

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US