

# Product datasheet for TP720850M

## NFYA (NM\_021705) Human Recombinant Protein

### **Product data:**

#### **Product Type: Recombinant Proteins Description:** Purified recombinant protein of Human nuclear transcription factor Y, alpha (NFYA), transcript variant 2 Species: Human **Expression Host:** E. coli **Expression cDNA Clone** Met1-Ser318 or AA Sequence: N-GST Tag: Predicted MW: 60.58 kDa **Concentration:** lot specific **Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl Endotoxin: Endotoxin level is $< 0.1 \text{ ng/}\mu\text{g}$ of protein ( $< 1 \text{ EU/}\mu\text{g}$ ) **Reconstitution Method:** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Storage: Store at -80°C. Stable for at least 6 months from date of receipt under proper storage and handling Stability: conditions. NP 068351 RefSeq: Locus ID: 4800 UniProt ID: P23511, A0A024RD22 6149 RefSeq Size: Cytogenetics: 6p21.1 **RefSeq ORF:** 954 Synonyms: CBF-A; CBF-B; HAP2; NF-YA



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Summary:	The protein encoded by this gene is one subunit of a trimeric complex, forming a highly conserved transcription factor that binds to CCAAT motifs in the promoter regions in a variety of genes. Subunit A associates with a tight dimer composed of the B and C subunits, resulting in a trimer that binds to DNA with high specificity and affinity. The sequence specific interactions of the complex are made by the A subunit, suggesting a role as the regulatory subunit. In addition, there is evidence of post-transcriptional regulation in this gene product, either by protein degradation or control of translation. Further regulation is represented by alternative splicing in the glutamine-rich activation domain, with clear tissue-specific preferences for the two isoforms. [provided by RefSeq, Jul 2008]
Protein Families: Protein Pathway	

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US