

Product datasheet for **TP720869XL**

DCK (NM_000788) Human Recombinant Protein

Product data:

| | |
|--|---|
| Product Type: | Recombinant Proteins |
| Description: | Purified recombinant protein of Human deoxycytidine kinase (DCK) |
| Species: | Human |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | Met1-Leu260 |
| Tag: | N-His&T7 |
| Predicted MW: | 34 kDa |
| Purity: | >95% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | Provided lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl |
| Endotoxin: | Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg) |
| Storage: | Store at -80°C. |
| Stability: | Stable for at least 3 months from date of receipt under proper storage and handling conditions. |
| RefSeq: | NP_000779 |
| Locus ID: | 1633 |
| UniProt ID: | P27707 , F5CTF3 |
| RefSeq Size: | 2618 |
| Cytogenetics: | 4q13.3 |
| RefSeq ORF: | 780 |
| Summary: | Deoxycytidine kinase (DCK) is required for the phosphorylation of several deoxyribonucleosides and their nucleoside analogs. Deficiency of DCK is associated with resistance to antiviral and anticancer chemotherapeutic agents. Conversely, increased deoxycytidine kinase activity is associated with increased activation of these compounds to cytotoxic nucleoside triphosphate derivatives. DCK is clinically important because of its relationship to drug resistance and sensitivity. [provided by RefSeq, Jul 2008] |
| Protein Families: | Druggable Genome |



[View online »](#)

Protein Pathways: Purine metabolism, Pyrimidine metabolism